

wherein n is 0-3 and each X is independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{NO}_2$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{OR}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{R}^{5'}$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_1\text{-C}_{10}$ alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_6\text{-C}_{14}$ aryl, $\text{C}_7\text{-C}_{24}$ alkaryl, $\text{C}_3\text{-C}_{13}$ heteroaryl, $\text{C}_4\text{-C}_{23}$ alkheteroaryl, substituted $\text{C}_1\text{-C}_{10}$ alkyl, substituted $\text{C}_2\text{-C}_{10}$ alkenyl, substituted $\text{C}_1\text{-C}_{10}$ alkoxy, substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, substituted $\text{C}_4\text{-C}_{23}$ alkheteroaryl and $-\text{Y-Ar}$;

wherein if X is a substituted group, it is substituted by one or more substituents independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NR}^5\text{R}^{5'}$, $-\text{NO}_2$, $-\text{NR}^5\text{C}(\text{O})\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{OR}^{5'}$ and halogen up to per-halosubstitution;

wherein R^5 and $\text{R}^{5'}$ are independently selected from H, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_6\text{-C}_{14}$ aryl, $\text{C}_3\text{-C}_{13}$ heteroaryl, $\text{C}_7\text{-C}_{24}$ alkaryl, $\text{C}_4\text{-C}_{23}$ alkheteroaryl, up to per-halosubstituted $\text{C}_1\text{-C}_{10}$ alkyl, up to per-halosubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, up to per-halosubstituted $\text{C}_2\text{-C}_{10}$ alkenyl, up to per-halosubstituted $\text{C}_6\text{-C}_{14}$ aryl and up to per-halosubstituted $\text{C}_3\text{-C}_{13}$ heteroaryl,

wherein Y is $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^5)-$, $-(\text{CH}_2)_m-$, $-\text{C}(\text{O})-$, $-\text{CH}(\text{OH})-$, $-(\text{CH}_2)_m\text{O}-$, $-(\text{CH}_2)_m\text{S}-$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)-$, $-\text{O}(\text{CH}_2)_m-$, $-\text{CHX}^a$, $-\text{NR}^5\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^5-$, $-\text{CX}^a_{2-}$, $-\text{S}(\text{CH}_2)_m-$ and $-\text{N}(\text{R}^5)(\text{CH}_2)_m-$,

$m = 1-3$, and X^a is halogen; and


Ar is a 5-10 member aromatic structure containing 0-4 members of the group consisting of nitrogen, oxygen and sulfur which is unsubstituted or substituted by halogen up to per-halosubstitution and optionally substituted by Z_{n1} ,

wherein $n1$ is 0 to 3 and each Z is independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{C}(\text{O})-\text{NR}^5$, $-\text{NO}_2$, $=\text{O}$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NR}^5\text{R}^{5'}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{SO}_2\text{R}^5$, $-\text{SO}_2\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{OR}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{R}^{5'}$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_1\text{-C}_{10}$ alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_6\text{-C}_{14}$ aryl, $\text{C}_3\text{-C}_{13}$ heteroaryl, $\text{C}_7\text{-C}_{24}$ alkaryl, $\text{C}_4\text{-C}_{23}$ alkheteroaryl, substituted $\text{C}_1\text{-C}_{10}$ alkyl, substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, substituted $\text{C}_7\text{-C}_{24}$ alkaryl and substituted $\text{C}_4\text{-C}_{23}$ alkheteroaryl;

wherein if Z is a substituted group, it is substituted by the one or more substituents independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $=\text{O}$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NO}_2$, $-\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{R}^{5'}$,

$-\text{NR}^5\text{C}(\text{O})\text{OR}^{5'}$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_1\text{-C}_{10}$ alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, C-C_{10} heteroaryl, $\text{C}_6\text{-C}_{14}$ aryl, $\text{C}_4\text{-C}_{24}$ alkheteroaryl and $\text{C}_7\text{-C}_{24}$ alkaryl

A is a heteroaryl moiety selected from the group consisting of



wherein

R^1 is selected from the group consisting of halogen, $\text{C}_3\text{-C}_{10}$ alkyl, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_1\text{-C}_{13}$ heteroaryl, $\text{C}_6\text{-C}_{14}$ aryl, $\text{C}_7\text{-C}_{24}$ alkaryl, up to per-halosubstituted $\text{C}_1\text{-C}_{10}$ alkyl, up to per-halosubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, up to per-halosubstituted $\text{C}_1\text{-C}_{13}$ heteroaryl, up to per-halosubstituted $\text{C}_6\text{-C}_{14}$ aryl, and up to per-halosubstituted $\text{C}_7\text{-C}_{24}$ alkaryl;

R^2 is selected from the group consisting of H, $-\text{C}(\text{O})\text{R}^4$, $-\text{CO}_2\text{R}^4$, $-\text{C}(\text{O})\text{NR}^3\text{R}^{3'}$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_3\text{-C}_{10}$ cycloalkyl, $\text{C}_7\text{-C}_{24}$ alkaryl, $\text{C}_4\text{-C}_{23}$ alkheteroaryl, substituted $\text{C}_1\text{-C}_{10}$ alkyl, substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, substituted $\text{C}_7\text{-C}_{24}$ alkaryl and substituted $\text{C}_4\text{-C}_{23}$ alkheteroaryl,

where R^2 is a substituted group, it is substituted by one or more substituents independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^4$, $-\text{C}(\text{O})\text{-NR}^3\text{R}^{3'}$,

-NO₂, -OR⁴, -SR⁴, and halogen up to per-halosubstitution,

wherein R³ and R^{3'} are independently selected from the group consisting of H, -OR⁴, -SR⁴, -NR⁴R^{4'}, -C(O)R⁴, -CO₂R⁴, -C(O)NR⁴R^{4'}, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₆-C₁₄ aryl, C₃-C₁₃ heteroaryl, C₇-C₂₄ alkaryl, C₄-C₂₃ alkheteroaryl, up to per-halosubstituted C₁-C₁₀ alkyl, up to per-halosubstituted C₃-C₁₀ cycloalkyl, up to per-halosubstituted C₆-C₁₄ aryl and up to per-halosubstituted C₃-C₁₃ heteroaryl; and


wherein R⁴ and R^{4'} are independently selected from the group consisting of H, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, C₆-C₁₄ aryl, C₃-C₁₃ heteroaryl, C₇-C₂₄ alkaryl, C₄-C₂₃ alkheteroaryl, up to per-halosubstituted C₁-C₁₀ alkyl, up to per-halosubstituted C₃-C₁₀ cycloalkyl, up to per-halosubstituted C₆-C₁₄ aryl and up to per-halosubstituted C₃-C₁₃ heteroaryl,

R^a is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, up to per-halosubstituted C₁-C₁₀ alkyl and up to per-halosubstituted C₃-C₁₀ cycloalkyl; and

R^b is hydrogen or halogen,

R^c is hydrogen, halogen, C₁-C₁₀ alkyl, up to per-halosubstituted C₁-C₁₀ alkyl or combines with R¹ and the ring carbon atoms to which R¹ and R^c are bound to form a 5- or 6-membered cycloalkyl, aryl or hetaryl ring with 0-2 members selected from O, N and S

3. (Amended) A method of claim 1, wherein B is




wherein Y is selected from the group consisting of -O-, -S-, -CH₂-, -SCH₂-, -CH₂S-, -CH(OH)-, -C(O)-, -CX^a₂, -CX^aH-, -CH₂O- and -OCH₂-, where X^a is halogen,

Q is a six member aromatic structure containing 0-2 nitrogen, substituted or unsubstituted by halogen, up to per-halosubstitution;

Q¹ is a mono- or bicyclic aromatic structure of 3 to 10 carbon atoms and 0-4 members of the group consisting of N, O and S, unsubstituted or unsubstituted by halogen up to per-halosubstitution, and


X, Z, n and n1 are as defined in claim 1 and s is 0 or 1.

6. (Amended) A method as in claim 5, wherein B is 2,3-dichlorophenyl or of the formula



wherein Q is phenyl, Q¹ is phenyl or pyridinyl, Y is -O-, -S-, -CH₂- or -SCH₂, X is CF₃, and Z is -OH, -Cl or NHC(O)-C_pH_{2p+1}, where p = 2-4, s = 0 or 1, n = 0 and n1 = 0 or 1.

14. (Amended) A method as in claim 13, wherein B is 2,3-dichlorophenyl or of the formula



wherein Q is phenyl, Q¹ is phenyl, pyridinyl or benzothiazolyl, Y is -O-, -S-, -CH₂- or -NH-, Z is Cl, -CH₃ or -OCH₃, s = 0 or 1, n = 0 and n1 = 0 or 1.

28. (Amended) A method as in claim 1, wherein the compound for formula I displays p38 IC₅₀'s of less than 10 μm as determined by an in-vitro p38 kinase inhibition assay.

29. (Amended) A method according to claim 1, wherein the disease is mediated by a cytokine and/or protease (proteolytic enzyme) regulated by p38.

31. (Amended) A method according to claim 29, comprising administering an amount of a compound of formula I effective to inhibit production of a disease-mediating cytokine or protease.

Please add new claims 42-53 as follows:

—42. (New) A method for the treatment of cancerous cell growth mediated by raf kinase comprising administering a compound of formula I.



wherein B is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, naphthyl, quinolinyl, isoquinolinyl, phthalimidinyl, furyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzofuryl, benzothienyl, indolyl, benzopyrazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl or benzisothiazolyl, substituted by one or more substituents independently selected from the group consisting of halogen, up to per-halosubstitution, and X_n , wherein n is 0-3 and each X is independently selected from the group consisting of $-\text{CN}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{NO}_2$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{OR}^5$, $-\text{NR}^5\text{C}(\text{O})\text{R}^5$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_1\text{-C}_{10}$ alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, phenyl, pyridinyl, naphthyl, isoquinolinyl, quinolinyl up to per halo-substituted $\text{C}_1\text{-C}_{10}$ alkyl, up to per halo-substituted $\text{C}_2\text{-C}_{10}$ alkenyl, up to per halo-substituted $\text{C}_1\text{-C}_{10}$ alkoxy, up to per halo-substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, and $-\text{Y-Ar}$;

wherein R^5 and $\text{R}^{5'}$ are independently selected from H, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_3\text{-C}_{10}$ cycloalkyl, up to per-halosubstituted $\text{C}_1\text{-C}_{10}$ alkyl, up to per-halosubstituted $\text{C}_2\text{-C}_{10}$ alkenyl and up to per-halosubstituted $\text{C}_3\text{-C}_{10}$ cycloalkyl,


wherein Y is $-\text{O}-$, $-\text{S}-$, $-\text{N}(\text{R}^5)-$, $-(\text{CH}_2)_m-$, $-\text{C}(\text{O})-$, $-\text{CH}(\text{OH})-$, $-(\text{CH}_2)_m\text{O}-$, $-\text{NR}^5\text{C}(\text{O})\text{NR}^5\text{NR}^{5'}$, $-\text{NR}^5\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{NR}^5-$, $-(\text{CH}_2)_m\text{S}-$, $-(\text{CH}_2)_m\text{N}(\text{R}^5)-$, $-\text{O}(\text{CH}_2)_m-$, $-\text{CHX}^a$, $-\text{CX}^a_2-$, $-\text{S}-(\text{CH}_2)_m-$ and $-\text{N}(\text{R}^5)(\text{CH}_2)_m-$,

$m = 1, 3$, and X^a is halogen; and

Ar is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, naphthyl, quinolinyl, isoquinolinyl, phthalimidinyl, furyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzofuryl, benzothienyl, indolyl, benzopyrazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl or benzisothiazolyl, optionally substituted by halogen up to per-halosubstitution and optionally substituted by Z_{n1} , wherein $n1$ is 0 to 3 and each Z is independently selected from the group consisting of $-\text{CN}$, $=\text{O}$, $-\text{CO}_2\text{R}^5$, $-\text{C}(\text{O})\text{NR}^5\text{R}^{5'}$, $-\text{C}(\text{O})\text{R}^5$, $-\text{NO}_2$, $-\text{OR}^5$, $-\text{SR}^5$, $-\text{NR}^5\text{R}^{5'}$, $-\text{NR}^5\text{C}(\text{O})\text{OR}^5$, $-\text{NR}^5\text{C}(\text{O})\text{R}^5$, $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_2\text{-C}_{10}$ alkenyl, $\text{C}_1\text{-C}_{10}$ alkoxy, $\text{C}_3\text{-C}_{10}$ cycloalkyl, phenyl, pyridinyl, naphthyl, isoquinolinyl, quinolinyl up to per halo-substituted $\text{C}_1\text{-C}_{10}$ alkyl, up to per halo-substituted $\text{C}_2\text{-C}_{10}$ alkenyl, up to per halo-substituted $\text{C}_1\text{-C}_{10}$ alkoxy, up to per halo-substituted $\text{C}_3\text{-C}_{10}$ cycloalkyl, and $-\text{Y-Ar}$;

$C(O)NR^5R^{5'}$, $-C(O)-NR^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-NR^5R^{5'}$, $-NR^5C(O)OR^{5'}$, $-C(O)R^5$, $-NR^5C(O)R^{5'}$, $-SO_2R^5$, $SO_2NR^5R^{5'}$, C_1-C_{10} alkyl, C_1-C_{10} alkoxy, C_3-C_{10} cycloalkyl, up to per halo-substituted C_1-C_{10} alkyl, and up to per halo-substituted C_3-C_{10} cycloalkyl, and

A is a heteroaryl moiety selected from the group consisting of



wherein

R^1 is selected from the group consisting of halogen, C_3-C_{10} alkyl, C_3-C_{10} cycloalkyl, C_1-C_{13} heteroaryl, C_6-14 aryl, C_7-24 alkaryl, up to per-halosubstituted C_1-C_{10} alkyl, up to per-halosubstituted C_3-C_{10} cycloalkyl, up to per-halosubstituted C_1-C_{13} heteroaryl, up to per-halosubstituted C_6-14 aryl, and up to per-halosubstituted C_7-24 alkaryl;

R^2 is selected from the group consisting of H, $-C(O)R^4$, $-CO_2R^4$, $-C(O)NR^3R^{3'}$, C_1-C_{10} alkyl, C_3-C_{10} cycloalkyl, C_7-C_{24} alkaryl, C_4-C_{23} alkheteroaryl, substituted C_1-C_{10} alkyl, substituted C_3-C_{10} cycloalkyl, substituted C_7-C_{24} alkaryl and substituted C_4-C_{23} alkheteroaryl,

where R^2 is a substituted group, it is substituted by one or more substituents independently selected from the group consisting of $-CN$, $-CO_2R^4$, $-C(O)-NR^3R^{3'}$, $-NO_2$, $-OR^4$, $-SR^4$, and halogen up to per-halosubstitution,

wherein R^3 and $R^{3'}$ are independently selected from the group consisting of H, $-OR^4$, $-SR^4$, $-NR^4R^{4'}$, $-C(O)R^4$, $-CO_2R^4$, $-C(O)NR^4R^{4'}$, C_1-C_{10} alkyl, C_3-C_{10} cycloalkyl, phenyl, pyridinyl, naphthyl, isoquinolinyl or quinolinyl

up to per-halosubstituted C₁-C₁₀ alkyl, up to per-halosubstituted C₃-C₁₀ cycloalkyl, and up to per-halosubstituted, phenyl, pyridinyl, naphthyl, isoquinolinyl or quinolinyl and


wherein R⁴ and R^{4'} are independently selected from the group consisting of H, C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, , phenyl, pyridinyl, naphthyl, isoquinolinyl, quinolinyl up to per-halosubstituted C₁-C₁₀ alkyl, up to per-halosubstituted C₃-C₁₀ cycloalkyl, and up to per-halosubstituted, phenyl, pyridinyl, naphthyl, isoquinolinyl or quinolinyl,

R^a is C₁-C₁₀ alkyl, C₃-C₁₀ cycloalkyl, up to per-halosubstituted C₁-C₁₀ alkyl and up to per-halosubstituted C₃-C₁₀ cycloalkyl; and

R^b is hydrogen or halogen,

R^c is hydrogen, halogen, C₁-C₁₀ alkyl, up to per-halosubstituted C₁-C₁₀ alkyl or combines with R¹ and the ring carbon atoms to which R¹ and R^c are bound to form a 5- or 6-membered cycloalkyl, aryl or hetaryl ring with 0-2 members selected from O, N and S.

43. (New) A method as in claim 42, wherein B is



which is substituted or unsubstituted by halogen, up to per-halosubstitution, and wherein

n = 1-3 and


each X is independently selected from the group consisting of C₁₋₄ alkyl, up to per-halosubstituted C₁₋₄ alkyl and -Y-Ar;

wherein Y is - O-, -S-, -N(R⁵)-, -(CH₂)_m-, -C(O)-, -CH(OH)-, -(CH₂)_mO-, -NR⁵C(O)NR⁵NR^{5'}-, -NR⁵C(O)-, -C(O)NR⁵-, -(CH₂)_mS-, -(CH₂)_mN(R⁵)-, -O(CH₂)_m-, -CHX^a-, -CX^a₂-, -S-(CH₂)_m- and -N(R⁵)(CH₂)_m-,

m = 1-3, and X^a is halogen; and


Ar is phenyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, naphthyl, quinolinyl, isoquinolinyl, phthalimidinyl, furyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, benzofuryl, benzothienyl, indolyl, benzopyrazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl or benzisothiazolyl, optionally substituted by halogen up to per-halosubstitution and optionally substituted by Z_{n1} , wherein $n1$ is 0 to 3 and each Z is independently selected from the group consisting of $-CN$, $=O$, $-CO_2R^5$, $-C(O)NR^5R^{5'}$, $-C(O)R^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-NR^5R^{5'}$, $-NR^5C(O)OR^{5'}$, $-C(O)R^5$, $-NR^5C(O)R^{5'}$, $-SO_2R^5$, $-SO_2R^5R^{5'}$, C_1-C_{10} alkyl, C_1-C_{10} alkoxy, C_3-C_{10} cycloalkyl, up to per halo-substituted C_1-C_{10} alkyl, and up to per halo-substituted C_3-C_{10} cycloalkyl, wherein R^5 and $R^{5'}$ are independently selected from H, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_3-C_{10} cycloalkyl, up to per-halosubstituted C_1-C_{10} alkyl, up to per-halosubstituted C_2-C_{10} alkenyl and up to per-halosubstituted C_3-C_{10} cycloalkyl.

44. (New) A method as in claim 5, wherein B is of the formula




wherein Q is phenyl or pyridinyl, optionally substituted by halogen up to per-halosubstitution, Q^1 is pyridinyl, phenyl or benzothiazolyl optionally substituted by halogen up to per-halosubstitution, Y is $-O-$, $-S-$, $-CH_2S-$, $-SCH_2-$, $-CH_2O-$, $-OCH_2-$ or $-CH_2-$, X is C_1-C_4 alkyl or up to per-halosubstituted C_1-C_4 alkyl and Z is as defined in claim 1, $n = 0$ or 1 , $s = 1$ and $n1 = 0-1$.

45. (New) A method as in claim 9, wherein B is of the formula




Q is phenyl or pyridinyl, optionally substituted by halogen up to per-halosubstitution, Q¹ is pyridinyl, phenyl or benzothiazolyl optionally substituted by halogen up to per-halosubstitution, Y is -O-, -S-, -C(O)- or -CH₂-, X is C₁-C₄ alkyl or up to per-halosubstituted C₁-C₄ alkyl and Z is as defined in claim 1 n = 0 or 1, s = 0 or 1 and n1 = 0 or 1.

46. (New) A method as in claim 13, wherein B is of the formula




Q is phenyl or pyridinyl optionally substituted by halogen up to per-halosubstitution, Q¹ is phenyl, benzothiazolyl or pyridinyl optionally substituted by halogen up to per-halosubstitution, Y is -O-, -S- or -CH₂-, X is C₁-C₄ alkyl or up to per-halosubstituted C₁-C₄ alkyl, Z is as defined in claim 1, n = 0 or 1, s = 1, and n1 = 0 or 1.

47. (New) A method as in claim 17, wherein B is of the formula




wherein Q is phenyl optionally substituted by halogen up to per-halosubstitution, Q¹ is phenyl or pyridinyl optionally substituted by halogen up to per-halosubstitution, Y is -O- or -S-, X is C₁-C₄ alkyl or up to per-halosubstituted C₁-C₄ alkyl, Z is as defined in claim 1, n = 0 or 1, s = 0 or 1 and n1 = 0-2.

48. (New) A method as in claim 22, wherein B is of the formula



wherein Q is phenyl optionally substituted by halogen up to per-halosubstitution, Q¹ is phenyl or pyridinyl optionally substituted by halogen up to per-halosubstitution, Y is -O- or -S-, X is C₁-C₄ alkyl or up to per-halosubstituted C₁-C₄ alkyl, s = 1, Z is as defined in claim 1, n = 0 or 1 and n1 = 0 or 1.

49. (New) A method as in claim 28, wherein B is of the formula



wherein Q is phenyl optionally substituted by halogen up to per-halosubstitution, Q¹ is phenyl or pyridinyl optionally substituted by halogen up to per-halosubstitution, and Y is -O- or -S-, X is C₁-C₄ alkyl or up to per-halosubstituted C₁-C₄ alkyl, Z is as defined in claim 1, n = 0 or 1, s = 0 or 1 and n₁ = 0-2.

50. (New) A method as in claim 1, wherein B is

a) phenyl, pyridinyl, naphthyl, quinolinyl or isoquinolinyl, substituted by -Y-Ar and optionally substituted by

- halogen up to per-halosubstitution,
- C₁-C₄ alkyl,
- up to per-halosubstituted C₁-C₄ alkyl, or
- a combination thereof,

wherein Y and Ar are as defined in claim 1;

- b) thienyl substituted by methyl; or
- c) indolyl substituted by phenyl or pyridyl.


51. (NEW) A method as in claim 1, wherein B is phenyl or pyridinyl substituted by -Y-Ar and optionally substituted by

- halogen, up to per-halosubstitution,
- C₁-C₄ alkyl,
- up to per-halosubstituted C₁-C₄ alkyl, or
- a combination thereof,

wherein Y and Ar are as defined in claim 1.


52. (NEW) A compound of one of the formulae

a)




b)

*BT
cm*




wherein R^6 is -O-CH₂-phenyl, -NH-C(O)-O-t-butyl, -O-n-pentyl, -O-n-butyl, -C(O)-N(CH₃)₂, -O-CH₂CH(CH₃)₂ or -O-n-propyl;

c)




wherein R^1 is -CH₂-t-butyl;


d)




wherein R^2 is -CH₂CF₃, -C₂H₄-OH, -CH₂-(3-HOC₆H₄), -CH₂C(O)NHCH₃, -CH₂C(O)OC₂H₅, -C₂H₄CN, or




e)
DT
cm



f)




g)



or

h)



and pharmaceutically acceptable salts thereof.

53. (NEW) A pharmaceutical composition comprising a compound according to claim 52 or a pharmaceutically acceptable salt thereof and a physiologically acceptable carrier.--